

- ¹²⁴ Harrison, L. “Foley catheter beats cook cervical for labor induction”. *Medscape Medical News*. 2015.
- ¹²⁵ Jozwiak, M.; Bloemenkamp, K.W.M.; Kelly, A.J.; Mol, B.W.J.; Irión, O.; Bouvain, M. “Mechanical methods for induction of labour”. *Cochrane Database of Systematic Reviews* 2012, Issue 3. Art. No.: CD001233. DOI: 10.1002/14651858.CD001233.pub2
- ¹²⁶ Ayres-de-Campos, D.; Montenegro, N.; Rodrigues, T.; Ramalho, C. *Protocolos de Medicina Materno-Fetal*. 3.^a edição. Lidel. 2014.
Indução do Trabalho de Parto. Orientação nº 002/2015 de 19/01/2015. Direcção-Geral da Saúde.
- ¹²⁷ “Guidelines on labor induction”. American College of Obstetricians and Gynecologists. Julho de 2009.
“Induction of Labor”. SOGC Clinical Practice Guideline. JOGC. 2015.
- ¹²⁸ *Cytotec*, folheto informativo: informação para o doente. Infarmed. 2013.
- ¹²⁹ Cardoso, L. *O uso off-label de medicamentos*. Universidade Fernando Pessoa. 2014.
- ¹³⁰ “Inducing Labor”. NICE Guideline CG70. 2008.
- ¹³¹ “Inducing Labor”. Queensland Clinical Guideline. Setembro de 2011.
- ¹³² “Use of prostaglandins for cervical ripening prior to the induction of labour”. RANZCOG. 2015.
- ¹³³ “Declaração Universal em Bioética e Direitos Humanos”. Art.º 6. UNESCO. 2005.
- ¹³⁴ “Consentimento Informado, Esclarecido e Livre Dado por Escrito”. Norma 015/2013. Direcção-Geral da Saúde. 2015.
- ¹³⁵ Clausen, J.A.; Juhl, M.; Rydahl, E. “Quality assessment of patient leaflets on misoprostol-induced labour: Does written information adhere to international standards for patient involvement and informed consent?”. *BMJ Open* 2016; 6:e011333. DOI:10.1136/bmjopen-2016-011333.
Lenk, C.; Duttge, G. “Ethical and legal framework and regulation for off-label use: European perspective”. *Therapeutics and Clinical Risk Management*. 2014; 10:537-546. DOI: 10.2147/TCRM.S40232.

Molyneux, C.; Bogaert, P. “The need of informed consent in off-label use in Europe”. *Informa UK*, Ltd. 2010.

¹³⁶ Allen, R.; O’Brien, B.M. “Uses of misoprostol in obstetrics and gynecology”. *Reviews in Obstetrics and Gynecology*. 2009; 2(3):159-168.

¹³⁷ Alfirevic, Z.; Aflaifel, N.; Weeks, A. “Oral misoprostol for induction of labour”. *Cochrane Database of Systematic Reviews* 2014, Issue 6. Art. No.: CD001338. DOI: 10.1002/14651858.CD001338.pub3.

¹³⁸ Hofmeyr, G.J.; Gülmezoglu, A.M.; Pileggi, C. “Vaginal misoprostol for cervical ripening and induction of labour”. *Cochrane Database of Systematic Reviews* 2010, Issue 10. Art. No.: CD000941. DOI: 10.1002/14651858.CD000941.pub2.

Alfirevic, Z.; Aflaifel, N.; Weeks, A. “Oral misoprostol for induction of labour”. *Cochrane Database of Systematic Reviews* 2014, Issue 6. Art. No.: CD001338. DOI: 10.1002/14651858.CD001338.pub3.

¹³⁹ Alfirevic, Z.; Keeney, E.; Dowswell, T. *et al.* “Methods to induce labour: a systematic review, network meta-analysis and cost-effectiveness analysis”. *Bjog*. 2016; 123(9):1462-1470. DOI: 10.1111/1471-0528.13981.

¹⁴⁰ Agência Europeia do Medicamento. Procedimento n.º PSU-SA/00010353/201505. <http://www.ema.europa.eu>.

¹⁴¹ “Indução do Trabalho de Parto”. Orientação n.º 002/2015 de 19/01/2015. Direcção-Geral da Saúde.

¹⁴² Alfirevic, Z.; Aflaifel, N.; Weeks, A. “Oral misoprostol for induction of labour”. *Cochrane Database of Systematic Reviews* 2014, Issue 6. Art. No.: CD001338. DOI: 10.1002/14651858.CD001338.pub3.

¹⁴³ WHO Recommendations for Labour Induction. Organização Mundial de Saúde. 2011.

¹⁴⁴ “Induction of labour: misoprostol vaginal delivery system”. NICE advice [ESNM38]. NICE. 2014. MISODEL 200 microgrammes, dispositif de diffusion vaginal. Commission de la transparence. HAS - Direction de l’Evaluation Médicale, Economique et de Santé Publique. 2015.

¹⁴⁵ Ayres-de-Campos, D.; Montenegro, N.; Rodrigues, T.; Ramalho, C. *Protocolos de Medicina Materno-Fetal*. 3.ª edição. Lidel. 2014.

¹⁴⁶ Alfirevic, Z.; Aflaifel, N.; Weeks, A. “Oral misoprostol for induction of

labour”. Cochrane Database of Systematic Reviews 2014, Issue 6. Art. No.: CD001338. DOI: 10.1002/14651858.CD001338.pub3.

¹⁴⁷ Chen, W.; Xue, J.; Peprah, M.K.; Wen, S.W.; Walker, M.; Gao, Y.; Tang, Y. “A systematic review and network meta-analysis comparing the use of Foley catheters, misoprostol, and dinoprostone for cervical ripening in the induction of labour”. *BJOG* 2016; 123:346-354.

¹⁴⁸ Lenk, C.; Duttge, G. “Ethical and legal framework and regulation for off-label use: European perspective”. *Therapeutics and Clinical Risk Management*. 2014;10:537-546. DOI: 10.2147/TCRM.S40232.

Clausen, J.A.; Juhl, M.; Rydahl, E. “Quality assessment of patient leaflets on misoprostol-induced labour: does written information adhere to international standards for patient involvement and informed consent?”. *BMJ Open* 2016;6:e011333. DOI: 10.1136/bmjopen-2016-011333.

¹⁴⁹ Alfirevic, Z.; Kelly, A.J.; Dowswell, T. “Intravenous oxytocin alone for cervical ripening and induction of labour”. The Cochrane database of systematic reviews. 2009;(4):CD003246. DOI:10.1002/14651858.CD003246.pub2.

¹⁵⁰ Boie, S.; Velu, A.V.; Glavind, J.; Mol, B.W.J.; Ulbjerg, N.; Graaf, I. de; Bor, P.; Bakker, J.J.H. “Discontinuation of intravenous oxytocin in the active phase of induced labour (Protocol)”. Cochrane Database of Systematic Reviews 2016, Issue 7. Art. No.: CD012274. DOI: 10.1002/14651858.CD012274.

¹⁵¹ Tsimis, M.S.; Buckley, A.P.; Nero, D.; Laurio, A. “Oxytocin usage for labor induction or augmentation and adverse neonatal outcomes”. American College of Obstetricians and Gynecologists 61st Annual Clinical Meeting; Nova Orleães, LA. 2013.

¹⁵² Boie, S.; Velu, A.V.; Glavind, J.; Mol, B.W.J.; Ulbjerg, N.; Graaf, I. de; Bor, P.; Bakker, J.J.H. “Discontinuation of intravenous oxytocin in the active phase of induced labour (Protocol)”. Cochrane Database of Systematic Reviews 2016, Issue 7. Art. No.: CD012274. DOI: 10.1002/14651858.CD012274.

¹⁵³ Daniel-Spiegel, E.; Weiner, Z.; Ben-Shlomo, I.; Shalev, E. “For how long should oxytocin be continued during induction of labour?”. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2004; 111: 331-334. DOI: 10.1111/j.1471-0528.2004.00096.x.

¹⁵⁴ Bor, P.; Ledertoug, S.; Boie, S.; Knoblauch, N.O.; Stornes, I. “Continuation versus discontinuation of oxytocin infusion during the active phase of labour: A randomised controlled trial”. *BJOG* 2016; 123:129-135.